


Standardization sessions improved urologists' agreement and boosted AI consistency in uroflowmetry interpretation.

Plata M¹, Azuero G¹, Daza F¹, Gutierrez A¹, Garcia V¹, Rojas-Rivillas M¹, Gomez S², Rodriguez T², Cañar S², Avila P², Gonzalez S², Florez S², Giraldo L²

¹Department of Urology, Fundación Santa Fe de Bogotá
²Universidad de los Andes

Improving Uroflowmetry Interpretation: Effects of Standardization Sessions on Interobserver Agreement and AI Model Consistency

Background



This study aimed to **assess** the **impact** of **standardization** sessions on **interobserver agreement** in **uroflowmetry** and their effect on **AI performance** and consistency.

Methods

- Weekly standardization sessions to review key urodynamic parameters until consensus.
- 50 uroflowmetries re-evaluated by 3 urologists pre- and post-standardization.
- Agreement: intraobserver (Fleiss' κ).
- AI: decision-tree models trained pre and post-standardization (F1-score).
- p < 0.05 significant.




Results

Table 1. Interobserver agreement

	Pre-standarization	Post-standarization
	Kappa	
Uroflowmetry	0.5	0.772
Curve morphology	0.5	0.513

Figure 1. Impact of standardization on AI decision-tree models



AI model mean **F1-score**: increased from 0.62 to **0.67**

Implications

- Standardization **reduces** interobserver **variability**.
- Greater agreement among urologists enhances AI performance and consistency.
- This dual benefit underscores the **potential** of integrating **human expertise** with **AI** to advance the **reliability** and **standardization** of urodynamic study interpretation.