

## W31: Diagnosis and repair of acute 3rd and 4th degree anal sphincter tears

Workshop Chair: Abdul Sultan, United Kingdom

27 August 2013 14:00 - 18:00

Start	End	Topic	Speakers
14:00	14:10	Introduction - Obstetric anal sphincter injuries (OASIS)	<ul style="list-style-type: none"> <li>Abdul Sultan</li> </ul>
14:10	14:30	anal anatomy and physiology	<ul style="list-style-type: none"> <li>Ranee Thakar</li> </ul>
14:30	14:40	anal endosonography	<ul style="list-style-type: none"> <li>Ranee Thakar</li> </ul>
14:40	15:10	diagnosis of OASIS	<ul style="list-style-type: none"> <li>Abdul Sultan</li> </ul>
15:10	15:30	repair of OASIS	<ul style="list-style-type: none"> <li>Abdul Sultan</li> </ul>
15:30	16:00	Break	None
16:00	16:15	management of subsequent pregnancy	<ul style="list-style-type: none"> <li>Abdul Sultan</li> </ul>
16:15	17:00	Videos on diagnosis and repair	<ul style="list-style-type: none"> <li>Ranee Thakar</li> </ul>
17:00	18:00	Hands-on repair of anal sphincters	<ul style="list-style-type: none"> <li>Abdul Sultan</li> <li>Ranee Thakar</li> </ul>

### Aims of course/workshop

Aim:

To learn how to identify, repair and manage primary obstetric anal sphincter injuries.

Objectives: (delegates will also have the opportunity to discuss these topics on an individual basis)

- Understand the anatomy and physiology of the anal sphincter.
- Learn the technique and interpretation of endoanal ultrasound
- Recognise and classify anal sphincter injury
- Observe a live video on diagnosis and repair
- Have hands on experience of repairing pig anal sphincters
- Understand the dilemmas regarding prevention and management of subsequent pregnancies
- Labour Ward protocol
- Have insight into the complications of anal sphincter trauma
- Set up a perineal clinic
- Skills to run a perineal trauma course

# Diagnosis & Repair of 3rd & 4th degree Tears

ICS 2013 Barcelona

## Hands-on Workshop

Croydon (Mayday) University Hospital  
Croydon, Surrey, UK

[abdul.sultan@croydonhealth.nhs.uk](mailto:abdul.sultan@croydonhealth.nhs.uk)

[raneethakar@croydonhealth.nhs.uk](mailto:raneethakar@croydonhealth.nhs.uk)

[www.perineum.net](http://www.perineum.net)

### Chairpersons

**Mr Abdul H Sultan, MB.ChB, MD, FRCOG**

*Consultant Obstetrician & Gynaecologist*

**Miss Raneethakar, MB BS, MD MRCOG**

*Consultant Obstetrician & Urogynaecologist*

## Applied anatomy and physiology of the perineum and anorectum Ranee Thakar

### Anatomy of the anorectum (Fig 1)


The anorectum is the most distal part of the gastrointestinal tract and consists of two parts: the anal canal and rectum. The anal canal measures about 3.5 cms and lies below the anorectal junction formed by the puborectalis muscle. The striated external anal sphincter (EAS) is made up of three parts (subcutaneous, superficial and deep) and is inseparable from the puborectalis dorsally. The internal anal sphincter (IAS) is a thickened continuation of the circular smooth muscle of the rectum. It is separated from the EAS by the conjoint longitudinal coat which is a continuation of the longitudinal smooth muscle of the rectum.

#### EAS:

- Striated muscle in a state of tonic contraction
- Innervated by the Pudendal nerve
- Up to 30% of resting pressure.
- Most of the squeeze pressure.
- Contraction maintained for < 2 minutes
- Reflex contraction with sudden increase in intra-abdominal pressure
- Relaxes during straining
- Damage results in urge faecal incontinence

#### IAS:

- Smooth muscle
- Autonomic control
- Contributes up to 70% of resting pressure
- Damage results in passive soiling and flatus incontinence




# Diagnosis of obstetric anal sphincter injuries (OASIS)

Abdul H Sultan

- Until the advent of anal ultrasound, the development of anal incontinence was attributed largely to pelvic neuropathy.
- However prospective studies before and after childbirth have shown that up to one third of women sustain anal sphincter damage that is not recognised at delivery (Sultan AH et al 1993).
- Andrews et al (2006) performed a study in which 241 women having their first vaginal delivery had their perineum re-examined by an experienced research fellow and endoanal ultrasound was performed immediately after delivery and repeated 7 weeks postpartum. When OASIS were identified by the research fellow, the injuries were confirmed and repaired by the duty registrar or consultant. The prevalence of clinically diagnosed OASIS increased from 11% to 25% (n=59). Every clinically diagnosed injury was identified by postpartum endoanal ultrasound. At 7 weeks no *de novo* defects were identified by ultrasound. This study concluded that most if not all sphincter defects that have previously been designated as "occult" injuries were in fact injuries that should have been recognisable at delivery. It was alarming to find that 87% and 27% of OASIS were not identified by midwives and doctors respectively. Although it is likely that some of these would have been detected at the time of suturing the tear, it is of concern that clinical recognition of OASIS is suboptimal.
- This finding is not unique as Groom and Patterson found that the rate of third degree tears rose to 15% when all "2<sup>nd</sup> degree tears" were re-examined by a second experienced person.
- It has been shown that only 16% of doctors and 39% of midwives feel that they were trained adequately to identify OASIS (Sultan et al 1995).
- On the other hand it is possible that the sphincter tear had been recognised but classified as a second-degree tear. A questionnaire sent to all UK consultants (Fernando et al 2002) and trainees (Sultan et al 1995) confirmed that up to 40% are still classifying partial and even complete disruption of the sphincter as a second degree. The reason for this confusion is partly due to previous teachings (Sultan & Thakar 2002) and therefore for the sake of clarification and consistency Sultan (1999) proposed a comprehensive classification that is now accepted by RCOG (Greentop guideline 2007), NICE (Nice.org.uk) and the International Consultation on Incontinence (Norton et al 2002) (Fig 2):

Fig 2: Classification of OASIS (Sultan 2007 Springer)



## OASIS – Classification (See Fig 2)

Sultan AH, *Clinical Risk* 1999; RCOG *Green Top Guidelines* 2001; ICI 2002; NICE 2007

1st degree = vaginal epithelium

2nd degree = perineal muscles

3rd degree = anal sphincter

3a = <50% external sphincter thickness

3b = > 50% external sphincter thickness

3c = internal sphincter torn

4th degree = 3<sup>rd</sup> degree + anal epithelium torn

# Repair techniques of obstetric anal sphincter injuries (OASIS)

*Abdul Sultan*

## **Anal incontinence after primary repair of OASIS** *Sultan AH, Thakar R 2007*

35 studies in the last 25 years

- Anal incontinence mean 39% (range 15 to 61%)
- Faecal incontinence mean 14% (range 2-29%)

## **Internal sphincter defects** *Mahony R et al 2007*

- 500 consecutive OASIS
- Persistent IAS defect independently associated with severe anal incontinence. OR 5.1 (95% CI = 1.5 – 22.9)

## **Fecal incontinence after vaginal delivery** *Fenner DE et al AJOG 2003*

- 831 primips completed bowel questionnaire 6 months after delivery
- 20% sustained OASIS
- 30% OASIS vs 20% of controls had poor bowel control.
- Symptoms 10x higher in 4<sup>th</sup> degree tears

## **Immediate –vs- delayed repair** *Nordenstam J et al 2008*

- RCT of 161 women
- Team of 3 obstetricians and 3 colorectal surgeons
- At 12 months 40% reported any anal incontinence (17% flatus > 1 per week)
- No difference in outcome between immediate and delayed (8 to 12 hours) repair

***No justification in delaying repair until the next day.***

## **Delayed and early secondary anal sphincter repair**

*Soerensen MM et al 2008*

- 21 female patients and 21 controls
- Delayed primary repair (<72 hours postpartum)
- Early secondary repair (<14 days postpartum)
- Repaired by 2 senior obstetricians
- Mean follow up of 4 years
- No post-op complications and none needed colostomy
- No significant difference in QoL with 19 controls

***25% vs 5% of controls had faecal incontinence***

**Anal canal length & good outcome** *Hool GR et al DCR 1998*

- Secondary overlap sphincter repair (n=51)
- Mean follow-up = 16 months
- Post-operative anal canal length best predicted continence

**Secondary anal sphincter repair** *Engel AF et al 1994; Malouf AJ et al 2000*

- Prospective study (n= 55) of overlap repair.
- 80% success at 18 months
- 50% at 5 years (n=46)
- But one third had more than one repair

**overlap vs end-to-end primary repair** *Sultan AH et al 1999*

- Anal incontinence: reduced from 42% to 8% (flatus)
- External sphincter defects: reduced from 85% to 15%
- Technique or operator? randomised study needed

**End-to-end vs overlap RCT** *Fernando R et al 2004*

- 64 randomised
- At one year compared to the end-to-end repair, significantly fewer women with overlap EAS repair suffered faecal incontinence
- 9 of 15 who had 3c/4<sup>th</sup> degree tear had FU scans
- All 9 had intact IAS.
  
- 119 primips 3b tear
- Primary outcome = solid stool leakage at least once per week
- No significant difference

**End-to-end vs overlap RCT**

- Farrell SA et al 2012
- 3 year follow-up
- No significant difference

### Suture materials ([www.perineum.net](http://www.perineum.net))

- Anal Mucosa - Vicryl 3-0
- Internal Anal Sphincter - Mattress end-to-end PDS 3-0
- External Anal Sphincter - Mattress/Overlap PDS 3-0

### Suture material *Williams et al 2006*

- 112 women – 4x4 randomised study
- No difference in suture related morbidity between Vicryl and PDS
- But 70% were 3a tears and only 54% 12 month follow-up

### Operating Theatre

- Sterile environment
- Good lighting
- Good exposure
- Appropriate instrument tray, sutures
- Anaesthesia – spinal, epidural, General
- Assistance

### Antibiotic prophylaxis for OASIS *Duggal N et al 2008*

- Prospective placebo controlled RCT (n=147)
- Single IV dose of cephalosporin
- Perineal wound infection 8% vs 24% in placebo


### OASIS repair - recommended practice *Sultan AH, Thakar R 2007*

- Experienced obstetrician
- Repair in operating theatre
- Regional or general anaesthesia
- IV antibiotics
- EAS
  - End-to-end for 3a
  - End-to-end or overlap for full thickness and full length 3b
- IAS → End-to-end mattress
- Monofilament sutures (PDS) for the sphincter (Vicryl 2-0 can also be used)
- Rectal examination before and after repair
- Foleys catheter for 12 hours
- Lactulose 15mls bd for 7 to 10 days
- Clinic Follow up in 2 to 3 months

### Labour Ward Protocol

See website [www.perineum.net](http://www.perineum.net)

**Fig 3 External anal sphincter defect (between arrows) on 3d endoanal ultrasound**



## **Management of OASIS after subsequent pregnancy**

**Abdul Sultan, Rane Thakar**

### **Mode of delivery after OASIS**

Caesarean section or Vaginal delivery?

### **Recurrence risks with previous OASIS** *Peleg D et al 1999*

- Primips, cep, term, 3<sup>o</sup>/4<sup>o</sup> (n=704); Incidence = 19% (midline episiotomy)
- Recurrence rate = 12% vs 7 % if no previous OASIS (P=0.001)

### **Previous OASIS - is recurrence predictable?** *Harkin R et al 2003*

- Mediolateral episiotomy
- 2 of 45 (4.4%) in subsequent vaginal deliveries developed a repeat OASIS

### **Previous OASIS** *Poen AC et al 1998*

- 43 of 110 women studied
- Anal incontinence 56% -v- 34% in women with no subsequent delivery. (RR = 1.6, CI = 1.1-2.5)

### **Previous OASIS** *Sangalli MR et al 2000*

- 177 women 13 years FU
- Faecal Incontinence in 114 subsequent deliveries (3<sup>o</sup> tears = 2.5% ; 4<sup>o</sup> tears = 26.5%)

### **Can OASIS be prevented ?** Can only minimise the risk of OASIS

- Episiotomy
  - Restrictive vs Liberal
  - Mediolateral vs Midline
- Instrumental delivery
  - Vacuum vs Forceps

### **Performing mediolateral episiotomy** *Andrews et al 2004; Andrews et al 2006*

- 254 primips, 41% mediolateral episiotomy
- No midwife and only 13 (22%) doctors performed a truly mediolateral episiotomy (between 40 to 60 degrees from the midline)
- Episiotomies angled closer to the midline significantly associated with OASIS (26 vs 37 degrees)

### **Episiotomy** *Eogan et al 2006*

- 50% risk reduction of third degree tears for every 6<sup>o</sup> away from midline

## Management of pregnancy after OASIS

Offer CS		Vaginal delivery
Asymptomatic	Hospital follow-up [Perineal Clinic]	Asymptomatic
Abnormal – ( Defect >1hr Incremental MSP < 20 mmHg)	Anal manometry & ultrasound	Normal
	Symptomatic	
severe faecal incontinence	Conservative Mx - Dietary advice - Regulate bowel action - Constipating agents: - codeine phosphate - loperamide - PFE & biofeedback	mild incontinence (flatus, staining)
Family not complete	Family complete/ delay	
Vaginal delivery	2° sphincter repair	Offer CS


For pictures, DVD, bibliography  
and suggested reading

See Website: [www.perineum.net](http://www.perineum.net)

2009. XII, 196 p. 82 illus., 32 in color. Softcover



ISBN 978-1-84800-996-7



## ***Take Home Messages***

---

- Every woman who has a vaginal delivery has a 3<sup>rd</sup> or 4<sup>th</sup> degree tear until proved otherwise
- A 3<sup>rd</sup> or 4<sup>th</sup> degree tear (as well as an isolated buttonhole tear of the rectum) cannot be excluded without a rectal examination
- Therefore a careful digital rectal examination with good exposure, adequate lighting and analgesia is mandatory.
- Restoration of normal sphincter length best predicts continence and therefore the full length of the torn sphincter must be repaired
- The torn internal sphincter should be identified and repaired separately using an end-to-end repair technique
- Partially torn external sphincter tears must be repaired using the end-to-end technique.
- Full thickness and full length external sphincter tears can be repaired by either the overlap or end-to-end technique. However the superiority of the overlap repair has only been demonstrated in one study where all repairs were conducted by only two operators. The Cochrane review does not recommend one technique over the other.
- The best chance of successful repair of OASIS is at the time of delivery; secondary sphincter repair (especially of the internal sphincter) is comparatively poor
- In terms of subsequent pregnancy management:
  - Compare like with like ie. the worst scenario of CS and VD
  - Risks following CS is not just for one CS but cumulative with each subsequent CS
  - Outcome of vaginal delivery after OASIS is good in selected patients

